





BIOS MeatFluid N 8,5 Organic nitrogen fluid fertilizers -

Fleshings fluid in suspension

Nitrogen

Nitrogen

Organic Carbon

N	Р	K
8,5	0	0

C	U.F.	
29,0	37,5	

Biological

QUICK TRANSFER - HYDROLISED PRODUCT (43% TOTAL AMINO ACIDS)

(N) total

(N) organic

(C) biological origin

BIOS MeatFluid N 8,5 is a nitrogenous organic fluid fertilizer, with high and balanced level of amino acids.

BIOS MeatFluid N 8,5 is "quickly absorption" fertilizer and it is useful for the needs of the plant in proteins, enzymes, growth regulators, sugars, etc.

BIOS MeatFluid N 8,5, applying it constantly, it gives an uniform vegetative development and improve quality and yeld of agricultural products, promoting their conservability; against stress (frosts, hail weeding, radical asphyxia), it helps the crops in all kinds of vegetative growth interruptions.

Finally, thanks to its wetting characteristics, anti-evaporate action and carrier functions, the use of BIOS MeatFluid N 8,5 increases the efficiency of the use of plant protection products.

COMPOSITION

P/P

8,5 %

8,5 %

29 %

equivalent to P/V at 20°C

10,6 %

10,6 %

36,3 %

Sill Sill Sill Sill Sill Sill Sill Sill	Z Z	
RICHEZZA RICHEZZA	BIOS MeatFluid N 8,5	

PACKAGING		
(Bottle Pz. 20 x Kg.1)		
Kg. 5 - 10 - 30 - 1.250		
Fluid		

Storage temperature: >=4° <=35° The product isn't combustible, and it's stable at ordinary temperature and pressure

	. J. De 6		400
	Carlo Carlo		
在 為了		1	
			100000

RAW MATERIALS	ORGANIC COMPONENT
NAW WATERIALS	Hydrolysed proteins, Residues of manufacturing of meat

RECOMMENDED DOSES - Kg./Ha				
Fertigation (quantity/application)	Kg./Ha			
rerugation (quantity/application)			rain	
Tree crops	30	-	60	
Horticulturar crops (open field)		-	60	
Greenhouses	30	-	90	
Foliar (quantity/intervention)				
from Kg. 1 a Kg. 6 for subministration/Ha				
Insert the diluted product since the first irrigation				

The recommended doses are indicative, must be increased and / or decreased taking into account: the climatic characteristics of the area of interest, fertility, water retention, soil structure, crop variety, the equipment used and the experience of the entrepreneur agricultural. In any case it is recommended to avoid concentrations of the product near the seed and / or roots



Restrictions on use - Article 11, paragraph 1, letter c) of Reg. EC / 1069/2009: The feeding of farm animals with herbaceous plants, taken through pasture or administered after being harvested, coming from land on which organic fertilizers or soil improvers other than manure have been applied, unless grazing or cutting of the grass takes place at the expiry of a waiting period of at least 21 days to ensure adequate control of health risks for public and animals.

The analytical data indicated on the packages follow the provisions of the Legislative Decree no. 75 of 29/04/2010 and subsequent amendments and / or additions. All data reported in this publication are indicative, BIOS s.r.l. reserves the right to modify them without prior notice.

KAKAKAKAKA